

## Five examples of Post Test Questions

---

1. During opening of the mouth, the arthrokinematic movement of rotation of the TMJ occurs:
  - A. in the upper joint space at the onset of opening
  - B. in the upper joint space only after 11mm of opening
  - C. in the lower joint space at the onset of opening
  - D. in the lower joint space only after 11mm of opening
  
2. Which of the following tissues contributes to the reduction of the disc on opening?
  - A. tension in the lateral pterygoid
  - B. tension in the TMJ ligament
  - C. tension in the posterior band
  - D. tension in the superior lamina
  
3. A clinical presentation of a disc displacement without reduction of the right TMJ is:
  - A. mouth opening is  $\leq 30$ mm with deviation to the right on opening
  - B. mouth opening is  $\leq 30$ mm with deflection to right on opening
  - C. mouth opening is  $\leq 30$ mm with deviation to the left on opening
  - D. mouth opening is  $\leq 30$ mm with deflection to left on opening
  
4. Which procedure relocates a disc displacement to a more normal position
  - a. TMJ Arthrocentesis
  - b. TMJ Arthroscopy
  - c. TMJ Arthrotomy
  - d. Orthognathic surgery
  
5. A possible adverse side effect from wearing a Nociceptive Trigeminal Inhibition (NTI) appliance is;
  - A. extrusion of posterior teeth
  - B. intrusion of posterior teeth
  - C. extrusion of anterior teeth
  - D. intrusion of anterior teeth